



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

*Death rate of Berlin and other cities.*

The death rate of Berlin for the week ended September 9 was lower than it has been for 5 weeks, amounting, calculated on the year, to 18.6 per thousand of the population, this being, however, higher than the rate for the corresponding week of last year, which amounted to only 17.7. Of the large towns and cities of Germany one-half showed less favorable health conditions than Berlin, including Königsberg, Breslau, Dresden, Leipzig, Halle, Magdeburg, Bremen, Cologne, Nuremberg, Stuttgart, Rixdorf (with 25.9). On the other hand, the following cities had a lower death rate than Berlin, namely: Hamburg, Munich, Carlsruhe, Frankfort on the Main, Cassel, Brunswick, Charlottenburg (with 16.2), Schöneberg (with 11.5), as well as London, Paris, and Vienna. The number of deaths among children in the first year of life showed again a decrease from the preceding week, while there was an increased mortality among the higher age classes. The infant death rate of Berlin, amounting to 7.2 per year and thousand, was higher than the Hamburg rate but lower than the Munich and Leipzig figures. Diseases of the digestive organs showed a considerable decrease, while acute diseases of the respiratory organs and infectious diseases remained almost stationary. There were registered 206 deaths from cholera (including 185 infants), 102 deaths from phthisis pulmonalis, 55 deaths from cancer, 50 deaths from acute diseases of the respiratory organs, 8 deaths from measles, 6 deaths from scarlet fever, 5 deaths from enteric fever, 3 deaths from diphtheria, and 3 deaths from influenza. Finally, 19 persons died by violence.

## GUATEMALA.

*Report from Livingston, fruit port.*

Acting Assistant Surgeon Peters reports as follows: Week ended September 23, 1905. Present officially estimated population, 3,500; 1 death; prevailing disease, malarial fever; general sanitary condition of this port and the surrounding country during the week, fair.

Bills of health were issued to the following-named vessels:

Date.	Vessel.	Destination.	Number of crew.	Number of passengers from this port.	Number of passengers in transit.	Pieces of baggage disinfected.
Sept. 19	Georg Dumois.....	New Orleans via Puerto Cortez.	19			
21	Belize .....	Mobile via Belize.	20			

*Reports from Guatemala City—Yellow fever along line of Northern Railroad.*

The following is received from Chargé d'Affaires ad interim Brown, under date of September 2.

Yellow fever, or a disease closely akin to it, has made disastrous progress during the past few months in the lowlands on the Atlantic slope of this country. It is difficult to ascertain with certainty all the facts. There seems to be no doubt, however, that there have been